



HSPG2 gene

heparan sulfate proteoglycan 2

Normal Function

The *HSPG2* gene provides instructions for making a protein called perlecan. This protein is found in the extracellular matrix, which is the intricate lattice of proteins and other molecules that forms in the spaces between cells. Specifically, it is found in part of the extracellular matrix called the basement membrane, which is a thin, sheet-like structure that separates and supports cells in many tissues. Perlecan is also found in cartilage, a tough, flexible tissue that makes up much of the skeleton during early development. Most cartilage is later converted to bone, except for the cartilage that continues to cover and protect the ends of bones and is present in the nose and external ears.

Perlecan is a heparan sulfate proteoglycan, which is a type of protein that interacts with many other proteins and has a variety of functions. In particular, perlecan is involved in cell signaling, the sticking (adhesion) of cells to one another, the formation of new blood vessels (angiogenesis), and the maintenance of basement membranes and cartilage throughout life. The protein also plays a critical role at the neuromuscular junction, which is the area between the ends of nerve cells and muscle cells where signals are relayed to trigger muscle contraction.

Health Conditions Related to Genetic Changes

Schwartz-Jampel syndrome

More than 30 mutations in the *HSPG2* gene have been found to cause Schwartz-Jampel syndrome. This rare condition is characterized by continuous muscle contraction (myotonia) that restricts movement, as well as bone abnormalities known as chondrodysplasia. Most of the mutations reduce the amount of perlecan that is produced. Other mutations lead to a version of perlecan that is only partially functional. A reduction in the amount or function of this protein disrupts the normal development of cartilage and bone tissue, which underlies chondrodysplasia in affected individuals. A reduced amount of functional perlecan at the neuromuscular junction likely alters the balance of other molecules that signal when muscles should contract and when they should relax. As a result, muscle contraction is triggered continuously, leading to myotonia.


other disorders

Mutations in the *HSPG2* gene can also cause another, more severe form of chondrodysplasia called dyssegmental dysplasia, Silverman-Handmaker type. Because of the very severe abnormalities associated with this rare condition, most affected individuals die before birth, are stillborn, or live only into early infancy. At least seven *HSPG2* gene mutations have been identified in people with dyssegmental dysplasia, Silverman-Handmaker type. These mutations prevent the production or transport of any functional perlecan. A total lack of this protein significantly disrupts the development of cartilage and bone tissue, causing this very severe type of chondrodysplasia.

Chromosomal Location

Cytogenetic Location: 1p36.12, which is the short (p) arm of chromosome 1 at position 36.12

Molecular Location: base pairs 21,822,232 to 21,937,297 on chromosome 1 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- endorepellin (domain V region)
- perlecan
- perlecan proteoglycan
- PLC
- PRCAN

Additional Information & Resources

Educational Resources

- Essentials of Glycobiology (second edition, 2009): Proteoglycans and Sulfated Glycosaminoglycans
<https://www.ncbi.nlm.nih.gov/books/NBK1900/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28HSPG2%5BTI%5D%29+OR+%28heparan+sulfate+proteoglycan+2%5BTI%5D%29+OR+%28perlecan%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>

OMIM

- DYSSEGMENTAL DYSPLASIA, SILVERMAN-HANDMAKER TYPE
<http://omim.org/entry/224410>
- HEPARAN SULFATE PROTEOGLYCAN OF BASEMENT MEMBRANE
<http://omim.org/entry/142461>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/HSPG2ID40890ch1p36.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=HSPG2%5Bgene%5D>
- HGNC Gene Family: I-set domain containing
<http://www.genenames.org/cgi-bin/genefamilies/set/593>
- HGNC Gene Family: Immunoglobulin like domain containing
<http://www.genenames.org/cgi-bin/genefamilies/set/594>
- HGNC Gene Family: Proteoglycans
<http://www.genenames.org/cgi-bin/genefamilies/set/570>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=5273
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/3339>
- UniProt
<http://www.uniprot.org/uniprot/P98160>

Sources for This Summary

- Arikawa-Hirasawa E, Le AH, Nishino I, Nonaka I, Ho NC, Francomano CA, Govindraj P, Hassell JR, Devaney JM, Spranger J, Stevenson RE, Iannaccone S, Dalakas MC, Yamada Y. Structural and functional mutations of the perlecan gene cause Schwartz-Jampel syndrome, with myotonic myopathy and chondrodysplasia. *Am J Hum Genet.* 2002 May;70(5):1368-75. Epub 2002 Apr 8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11941538>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC447613/>
- Arikawa-Hirasawa E, Wilcox WR, Le AH, Silverman N, Govindraj P, Hassell JR, Yamada Y. Dyssegmental dysplasia, Silverman-Handmaker type, is caused by functional null mutations of the perlecan gene. *Nat Genet.* 2001 Apr;27(4):431-4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11279527>
- Arikawa-Hirasawa E, Wilcox WR, Yamada Y. Dyssegmental dysplasia, Silverman-Handmaker type: unexpected role of perlecan in cartilage development. *Am J Med Genet.* 2001 Winter;106(4):254-7. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11891676>
- OMIM: HEPARAN SULFATE PROTEOGLYCAN OF BASEMENT MEMBRANE
<http://omim.org/entry/142461>
- Ladhani NN, Chitayat D, Nezarati MM, Laureane MC, Keating S, Silver RJ, Unger S, Velsher L, Sirkin W, Toi A, Glanc P. Dyssegmental dysplasia, Silverman-Handmaker type: prenatal ultrasound findings and molecular analysis. *Prenat Diagn.* 2013 Nov;33(11):1039-43. doi: 10.1002/pd.4193. Epub 2013 Aug 4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23836246>
- Melrose J, Hayes AJ, Whitelock JM, Little CB. Perlecan, the "jack of all trades" proteoglycan of cartilaginous weight-bearing connective tissues. *Bioessays.* 2008 May;30(5):457-69. doi: 10.1002/bies.20748. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18404701>
- Nicole S, Davoine CS, Topaloglu H, Cattolico L, Barral D, Beighton P, Hamida CB, Hammouda H, Cruaud C, White PS, Samson D, Urtizbereja JA, Lehmann-Horn F, Weissenbach J, Hentati F, Fontaine B. Perlecan, the major proteoglycan of basement membranes, is altered in patients with Schwartz-Jampel syndrome (chondrodystrophic myotonia). *Nat Genet.* 2000 Dec;26(4):480-3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11101850>
- Stum M, Davoine CS, Fontaine B, Nicole S. Schwartz-Jampel syndrome and perlecan deficiency. *Acta Myol.* 2005 Oct;24(2):89-92. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16550923>
- Stum M, Davoine CS, Vicart S, Guillot-Noël L, Topaloglu H, Carod-Artal FJ, Kayserili H, Hentati F, Merlini L, Urtizbereja JA, Hammouda el-H, Quan PC, Fontaine B, Nicole S. Spectrum of HSPG2 (Perlecan) mutations in patients with Schwartz-Jampel syndrome. *Hum Mutat.* 2006 Nov;27(11):1082-91.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16927315>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/HSPG2>

Reviewed: April 2016

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services